The Cosmological Legacy of Planck



Martin White **Berkeley.**

Figs. courtesy V. Pettorino

CMB encodes valuable information

- The CMB spectrum depends upon the initial spectrum of perturbations (inflation?) and the conditions in the photon-baryon fluid prior to last scattering.
- The rich structure in the spectrum, and the dependence on many cosmological parameters, provides a gold-mine of information.
- Scattering of an anisotropic temperature field generates (linear) polarization, which allows access to even more information.
- We can also get information about the low z Universe by looking at CMB lensing (and BAO – the sound waves frozen in the matter perturbations).

The magic of CMB ...

The CMB contains a gold-mine of information

- *if* it can be <u>accurately measured</u>
- and compared to precise theoretical predictions with a rich phenomenology
- in a statistically reliable
- and <u>computationally tractable</u> way

There are very few situations in cosmology, astrophysics (or indeed physics) where all of these conditions are met.

It is the intersection of these qualities that makes CMB such a powerful cosmological probe!

Planck mission

- Planck was a 3rd generation space mission (COBE, WMAP)
 - Like WMAP, Planck observed at "L2".
- It was part of ESA's "Cosmic Visions" program.
- It was the first sub-mm mission to map the entire sky to sub-Jy sensitivity and resolution better than 10 arcmins.
 - 74 detectors covering 25GHz-1000GHz, resolution 33'-5'.









CMB map: smoothed + polarization



Power spectrum ...

Hi, Dr. Elizabeth? Yeah, Uh... I accidentally took the Fourier transform of my cat... Meow! xkcd.com/26

The angular power spectrum



Polarization-Temperature



Polarization and lensing



Data compression!

- We find that a simple, 6 parameter model fits the data extremely well.
 - Data compression: trillions of bits of data are compressed to billions of measurements at 9 frequencies, then tens of millions of modes are compressed to thousands of multipoles which are compressed to 6 cosmological parameters!
 - With no evidence for a 7^{th} .
- For the "base model" the CMB determines all of the parameters, on its own, with exceptional accuracy.
 - If we include polarization, best determined parameter is 0.03%.
 - Only 1 parameter not determined to better than 1%.

Planck(-only) base ACDM model

Parameter	Description	Value	
ω_{b}	Baryon density	0.02237 ± 0.00015	
ω_{c}	Cold dark matter density	0.1200 ± 0.0012	
1000 _{MC}	Angular size of acoustic scale	1.04092 ± 0.00031	
τ	Optical depth to Thomson scattering	0.0544 ± 0.0073	
In(10 ¹⁰ A _s)	Observed fluctuation amplitude	3.044 ± 0.014	
n _s	Slope of primordial power spectrum (spectral index)	0.9649 ± 0.0042	
H ₀ (km/s/Mpc)	Expansion rate of Universe	67.36 ± 0.54	
σ ₈	Amplitude of fluctuations in matter today	0.8111 ± 0.006	

And my favorite derived parameter: $k_{eq} = 0.01038 \pm 0.00008 \text{ Mpc}^{-1}$

Improvement in parameters in 15yr



Acoustic scale: known to 0.03%



Limits on primordial perturbations ...



T & E consistency



Can use EE,TE to predict TT assuming Λ CDM.

Can use TT,TE to predict EE.

Can use TT,EE to predict TE.

All looks good!

CMB lensing

- Photons from the CMB are deflected on their way to us by the potentials due to large-scale structure.
- The typical deflection is 2-3 arcmin but deflections are coherent over degrees.
 - Signal dominated by structures of tens of Mpc at $z\sim 2$.
- Gives sensitivity to the "low z" Universe.
 - Allows us to break some degeneracies from purely within the CMB dataset.
 - Provides a cross-check on the paradigm: are the structures we infer at $z\sim2$ consistent with the "initial conditions" measured at $z\sim1,000$? [After 10³ growth: A_L^{\phi\phi}=0.997±0.03]
- Provides a map, over the whole sky, of the (projected) mass back to the surface of last-scattering (98% of the way to the horizon).

Lensing deflection (E-mode)



-0.0016

Lensing now measured at >40 σ . Better than predicted by anisotropy! Much future CMB science will be lensing ...

Some key "early Universe" results ...

- Inflation.
 - Planck has had a huge impact on inflationary model building!
 - A large number of "popular" models now ruled out.
 - The simplest models of inflation predict ...

A spatially flat Universe	$\Omega_{\rm K}$ =0.0007 ± 0.0019
with <i>nearly</i> scale-invariant (red) spectrum of density perturbations	0.967 ± 0.004
which is almost a power-law	dn _s /dln <i>k</i> = -0.0042 ± 0.0067
dominated by scalar perturbations	r _{0.002} <0.07 (95%)
which are Gaussian	f _{NL} = 2.5 ± 5.7 ~ 0
and adiabatic	α_{-1} = 0.00013 ± 0.00037
with negligible topological defects	f _{NG} < 0.01 (95%)

Inflationary models

- Coherence of peaks, sign of TE
 - Early Universe origin of perturbations
- $\Omega_{\rm K} \sim 0$: duration of slow-roll not fine tuned.
- Primordial P(k) well approximated by power-law.
 - Inflaton rolls down a featureless, nearly flat potential.
- No isocurvature modes: 1 d.o.f.
- Scalar modes dominate by 1 order of magnitude.
 - Models with $r\sim(1-n_s)$ severely limited.
 - Models with $r \sim (1-n_s)^2$ require next-gen technology to limit.
 - Models with $r < < (1-n_s)^2$ out of reach of foreseeable technology.
- Surviving models have V'~0 and V"<0</p>
 - special point in potential.

Some key "early Universe" results ...

Light neutrinos.

- Neutrinos non-relativistic at $z \sim 10^3$ long ruled out.
- Current constraints come primarily from lensing and distances.
- Starting to put pressure on inverted hierarchy.

Light relic species

- Dark, relativistic d.o.f. labeled by N_{eff}

$$\frac{\rho_{\rm rad}}{\rho_{\gamma}} = \frac{7}{8} \left(\frac{4}{11}\right)^{4/3} N_{\rm eff} \qquad N_{\rm eff} = 3.045 + \Delta N_{\rm eff}$$

Constraints on neutrinos now tighter $\Sigma m_v < 0.12 \text{ eV}$ (95%)





Light degrees of freedom: N_{eff}

CMB + LSS

- By cementing the gravitational instability paradigm and measuring the ICs and parameters, Planck sets the framework for LSS.
- Planck precisely determines many of the key parameters for large-scale structure:
 - $k_{eq} = 0.01038 \pm 0.00008 \text{ Mpc}^{-1}$
 - $\sigma_8(z=2) = 0.3211 \pm 0.0009$
 - $r_{drag} = 147.09 \pm 0.26$ Mpc
- Planck calibrates the "standard fluctuation spectrum".
 - Sets the scale and level of inhomogeneity in the Universe.
 - Governs structure formation, galaxy formation, etc.

Early on, the fields of LSS and CMB were tightly coupled. With time they grew apart and specialized. I think we are witnessing a re-coupling.

Low redshift structure



Rate of growth of large-scale structure



Redshift-space distortions: Planck vs BOSS

Growth rate ($f\sigma_8$) vs. A-P parameter (F_{AP})



(BOSS results have been marginalized over D_V: dashed lines show results conditioned on Planck)

Cosmic distance scale



Cosmic distance scale: SNe



Comparison with lensing



Just plain cool ...

- In 2013 Planck detected the motion of the Earth in the aberration of the measured CMB anisotropy.
 - Observed at >4 σ in 2013 data.
- In 2015 we detected the impact of fluctuations in the 2K neutrino background!
 - Evidence for v background strong (N_{eff} =0 ruled out @ >10 σ)
 - Now have exquisite detection of free-streaming of this component (measures of c_{eff}^2 and c_{vis}^2).
- In 2018 we measured the "gravitational slip" at z=1000 to be 1.004 ± 0.007.

- GR predicts it is 1.

Temperature story – begun by COBE – is (essentially) done ...



The next generation

- Search for polarization "B-modes"
 - Generated by primordial gravity waves
 - Constrains the energy scale of inflation.
- Primordial non-Gaussianity.
 - Details of inflationary dynamics.
- CMB lensing & cross-correlation.
 - Tests of gravity and large-scale modes.
 - Measurement of neutrino mass.
- tSZ and kSZ.
 - Probes of large-scale velocities, reionization & gastrophysics.

Conclusions

- The CMB is our premier cosmological laboratory.
- Experiments provide a rigorous test of our models using the physics of harmonic oscillators!
 - Established acoustic physics as the "gold standard" probe.
- Impressive confirmation of the standard cosmological model.
 - Precise constraints on model and parameters.
 - Tight limits on deviations from base model.
 - Some indications of internal and external tensions, but with only modest statistical significance.
- Next generation CMB experiments are underway, and planning for CMB-S4 is in progress ...
- Synergies between large-scale structure and CMB are only growing in importance!

Planck data

- All Planck papers can be downloaded from
 - http://www.cosmos.esa.int/web/planck/publications

Except

- > Power spectra, likelihood (& likelihood code).
- Isotropy and statistics
- > Primordial non-Gaussianity.
- All Planck data can be downloaded from
 - http://pla.esac.esa.int/pla





Changes: quick version

- For TT, very little change.
- For EE, tighter τ limit from low *I*
- EE and TE systematics reduced (but not eliminated roughly <0.5 σ left).
- $\phi\phi$ pushed to L=8 rather than 40.

You probably don't care, but the dipole amplitude is now known to 0.025% -- the same uncertainty as for the monopole (i.e. the temperature)!!!

Three CMB numbers known to <0.1%

Changes to ACDM parameters



Changes to ACDM parameters



Jackknife tests



Using statistical weight: σ⁻²



Inflation



Power law index of perturbation spectrum

Optical depth to Thomson scattering

New Planck results point to "late and fast" reionization.



Current Tensions: H₀

Even where tensions remain, dramatic progress has been made!



Current "tensions": clusters





World compilation of CMB anisotropy measurements

Planck dominates TT and (for now) the low ell TE and EE measurements. So far good consistency between experiments.

Physics is Universal!

Baryon density measured by BBN and CMB are in excellent agreement ... comparison uses all known laws of physics!



[And we also have a measurement of the Hydrogen 2s→1s transition which is 5x better than the lab measurement, and in fantastic agreement with the theoretical calculation!]

Consistency with other data

- The Planck data are consistent with the predictions of the simplest Λ CDM models.
- Within the framework of such models we can compare to a wide variety of other astrophysical/cosmological datasets.
 - Primordial nucleosynthesis
 - Baryon Acoustic Oscillations (distance scale).
 - Direct measures of H₀.
 - Redshift-space distortions.
 - Type la SNe.
 - Cosmic shear.
 - Counts of rich clusters of galaxies.
- Tensions remain.

- etc

Non-Gaussianity: f_{NL}

Туре	2013	2014	Generated by
Local	2.7±5.8	0.7±5.1	Curvaton, reheating, multifield,
Equilateral	-42±75	-9.5±44	Non-canonical kinetic term or higher derivative (e.g. K- flation, DBI, ghost inflation, with c _s <<1).
Orthogonal	-25±39	-25±22	Non-canonical kinetic term or higher derivative (c _s <<1).

(Other, specific shapes/cases are discussed in papers)



First there was COBE ...





Nobel prize in Physics, 2006, awarded to

Mather and Smoot

"for their discovery of the blackbody form and anisotropy of the cosmic microwave background radiation"







Ground based observations...



Angular power spectrum!



Anisotropy generates (linear) polarization



A quadrupole anisotropy generates linear poln.

Normally we define polarization patterns in terms of their parity and (confusingly!) refer to them as E & B modes.

Density perturbations can generate only E-mode polarization, but primordial gravity waves (or vorticity) can generate both E- and B-modes.

Baryon Acoustic Oscillations?

The oscillations in the photon-baryon fluid also imprint a feature in the latetime clustering of matter ... with the same characteristic length scale! Allows a "standard ruler" test of the expansion history!



CMB lensing & SZ effect(s)



"Secondary" anisotropies give us access to information in the "late" Universe – but require higher angular resolution and signal-to-noise ratio.





Looking back to the dawn of time



Planck Telescope 1.5x1.9m off-axis Gregorian T = 50 K





LFI Radiometers 30-70 GHz, T = 20 K



HFI Bolometers 100-857 GHz, T = 0.1 K



CMB anisotropy history

- Primordial anisotropy first detected by COBE in 1992.
 - Nobel prize to George F Smoot for "DMR".
- Ground and balloon borne experiments during the 1990s delineated the first peak and the damping tail and first measured polarization anisotropy.
- WMAP successor to COBE measured the first 2-3 peaks and begun to fill in the polarization story.
- Planck latest space mission currently provides our most precise measurements of temperature and polarization anisotropies.
 - Augmented at small scales by more sensitive, higher resolution, ground-based experiments.

The cosmic microwave background

- The entire Universe is filled with radiation in the form of a 2.7K black-body.
- This radiation is a relic of the hot, dense, early phase of the Universe (the hot-big bang).
- The light travels to us from a "surface of last scattering" at z~1100 (when the Universe was 10⁻³ times smaller than today and only 380,000yr old).
 - At this z the Universe was finally cold enough for protons to capture electrons to form neutral Hydrogen.
 - Optical depth to photon scattering quickly drops from τ >>1 to τ <<1.
- The radiation is almost the same intensity in all directions, but contains tiny fluctuations in intensity (or temperature) at the level of 10⁻⁴: CMB anisotropy.

The cartoon: sound waves in the early Universe

- At early times the universe was hot, dense and ionized. Photons and matter were tightly coupled by Thomson scattering.
 - Short m.f.p. allows fluid approximation.
- Initial fluctuations in density and gravitational potential drive acoustic waves in the bγ fluid: compressions and rarefactions.
- These show up as temperature fluctuations in the CMB, including an almost harmonic series of peaks in the angular power spectrum of ΔT as a function of angular wavenumber / (conjugate to angle θ).